



Medicinal Value of *Cynanchum Auriculatum* Plant: A Brief Study

Akhilesh Chandra Verma, Ph.D., Department of Chemistry
Government Naveen College Kui-Kukdur, Distt. Kabirdham, Chhattisgarh, INDIA

ORIGINAL ARTICLE



Author

Akhilesh Chandra Verma, Ph.D.

E-mail : dr.acverma8@gmail.com

shodhsamagam1@gmail.com

Received on : 05/03/2026
Revised on : 06/05/2026
Accepted on : 15/05/2026
Overall Similarity : 02% on 07/05/2026



Plagiarism Checker X - Report

Originality Assessment

2%

Overall Similarity

Date: May 7, 2026 (02:27 PM)
Matches: 74 / 3090 words
Sources: 3

Remarks: Low similarity detected, consider making necessary changes if needed.

Verify Report:
Scan this QR Code



ABSTRACT

The present study provides a systematic and critically integrated evaluation of the medicinal potential of *Cynanchum auriculatum*, a pharmacologically significant plant belonging to the family Asclepiadaceae. The analysis synthesizes phytochemical composition, biological activities, and ethnomedicinal relevance to establish a coherent understanding of its therapeutic value. Phytochemical investigations reveal the presence of structurally diverse bioactive constituents, including C21-steroids, acetophenones, alkaloids, terpenoids, and glycosides, which collectively underpin its pharmacodynamic properties [1], [2]. Among these, pregnane-type C21-steroids and specific compounds such as caudatin and cynandione A have demonstrated pronounced anti-tumor, hepatoprotective, neuroprotective, and anti-inflammatory activities [3]. The antioxidant efficacy of *C. auriculatum* is primarily attributed to its phenolic and flavonoid constituents, which facilitate efficient free radical scavenging and redox homeostasis [4]. In addition, its antimicrobial activity against a broad spectrum of bacterial and fungal strains underscores its potential as a natural therapeutic agent [5]. Ethnomedicinally, the plant has been extensively utilized in East Asian traditional systems for enhancing immunity, improving hepatic function, and managing chronic pathological conditions [6]. Despite its promising pharmacological profile, comprehensive validation in terms of pharmacokinetics, toxicity, and clinical efficacy remains limited. Therefore, this study emphasizes the necessity of integrating traditional knowledge systems with modern phytochemical and pharmacological approaches to establish *C. auriculatum* as a scientifically validated medicinal resource [7].

KEY WORDS

Cynanchum Auriculatum, *Phytochemical Analysis*, *Antioxidant Activity*, *Antimicrobial Activity*, *Ethnomedicine*, *Medicinal Plants*.

INTRODUCTION

Medicinal plants have historically constituted a foundational pillar of global healthcare systems, serving as a sustainable reservoir of bioactive compounds with diverse therapeutic applications. Their significance lies in the presence of structurally complex secondary metabolites capable of exerting multifaceted pharmacological effects with comparatively reduced adverse outcomes [1]. In recent decades, there has been a resurgence of scientific interest in plant-based therapeutics, particularly in the context of drug discovery and the management of chronic and degenerative diseases [2].

Ethnomedicine, encompassing traditional plant-based healing systems, continues to play a critical role in identifying biologically active species and guiding pharmacological research. A substantial proportion of modern pharmaceuticals have originated from ethnobotanical leads, thereby reinforcing the importance of preserving and scientifically validating traditional medicinal knowledge [6].

Cynanchum auriculatum Royle ex Wight, commonly referred to as “Baishouw” in traditional Chinese medicine, represents a therapeutically important species within the genus *Cynanchum*. It is widely distributed across East Asian regions, including China, Japan, and Korea, where it has been extensively employed for its medicinal properties [6]. However, its morphological and phytochemical resemblance to closely related species such as *Cynanchum bungei* and *Cynanchum wilfordii* often leads to challenges in accurate identification and standardization [7].

Pharmacological studies have demonstrated that *C. auriculatum* exhibits a broad spectrum of biological activities, including antioxidant, anti-inflammatory, hepatoprotective, immunomodulatory, and anti-tumor effects [3], [4]. These therapeutic attributes are primarily associated with its rich phytochemical profile, particularly the presence of C21-steroids and acetophenone derivatives [2].

Despite the growing body of literature, significant gaps persist in understanding its precise mechanisms of action, clinical efficacy and safety profile. Furthermore, issues related to species authentication and phytochemical standardization continue to hinder its translational application. Therefore, the present study aims to provide a structured and critical overview of *C. auriculatum*, focusing on its botanical characteristics, phytochemical composition, and medicinal significance, while highlighting future research directions [7].

Botanical and Phytochemical Overview

1. Taxonomy and Morphology

Cynanchum auriculatum is a perennial medicinal herb belonging to the family Asclepiadaceae and is taxonomically classified under the genus *Cynanchum*, which comprises numerous pharmacologically active species [1]. The plant exhibits distinctive morphological characteristics that facilitate its identification and differentiation from closely related taxa.

The roots are thick, cylindrical, and tuberous with a brownish-yellow outer surface, constituting the primary medicinal component. The stems are slender and often pubescent, while the leaves are oval to broadly elliptic, measuring approximately 4–12 cm in length and 4–10 cm in width. The plant produces small, white, oval-shaped flowers with a softly hairy internal surface, typically arranged in clustered inflorescences. The fruit is characterized as a double follicle, lanceolate in structure, containing seeds with filamentous appendages that facilitate dispersal [7].

The flowering period extends from June to September, followed by fruiting from July to November. These morphological attributes are critical for accurate botanical identification and play a vital role in minimizing misidentification, particularly in traditional herbal markets where closely related species coexist [6].

2. Phytochemical Basis

The therapeutic efficacy of *Cynanchum auriculatum* is intrinsically linked to its chemically diverse phytoconstituent profile. Extensive phytochemical investigations have revealed the presence of multiple classes

of secondary metabolites, with C21-steroids representing the dominant and pharmacologically significant group [2], [3].

In addition to C21-steroids, the plant contains acetophenones, terpenoids, alkaloids, glycosides, and other bioactive compounds that collectively contribute to its biological activity. Notably, compounds such as caudatin and cynandione A have been extensively studied for their therapeutic potential. C21-steroids, characterized by a pregnane-type structural framework, exhibit pronounced anti-tumor, antioxidant, and anti-inflammatory properties, whereas acetophenone derivatives contribute to hepatoprotective and neuroprotective effects [3].

The identification and characterization of these compounds are achieved through advanced analytical techniques, including high-performance liquid chromatography (HPLC), mass spectrometry (MS), nuclear magnetic resonance (NMR), and high-speed counter-current chromatography (HSCC). These methodologies enable precise structural elucidation and facilitate the correlation between chemical composition and pharmacological activity, thereby supporting targeted drug development strategies [2].

3. Medicinal Significance

The medicinal relevance of *Cynanchum auriculatum* is deeply rooted in traditional medical systems, particularly in East Asian ethnomedicine, where it has been employed for centuries in the management of various physiological disorders [6]. Traditionally, the plant has been used for nourishing blood, enhancing hepatic function, strengthening renal activity, and improving systemic vitality.

Modern pharmacological studies have substantiated these traditional claims by demonstrating its significant antioxidant, antimicrobial, anti-inflammatory, and immunomodulatory properties [4], [5]. The antioxidant activity plays a crucial role in mitigating oxidative stress through the neutralization of reactive oxygen species, thereby preventing cellular and molecular damage. Concurrently, its antimicrobial properties contribute to its efficacy against pathogenic microorganisms.

Furthermore, the plant has shown promising potential in the management of chronic diseases, including cancer, liver disorders, and metabolic dysfunctions, primarily due to its rich bioactive composition [3]. The convergence of traditional knowledge and contemporary scientific validation reinforces the therapeutic significance of *C. auriculatum*, positioning it as a promising candidate for further pharmacological investigation and clinical application [7].

Materials and Methods

1. Plant Collection and Authentication

The plant material of *Cynanchum auriculatum* was collected from an ecologically suitable and geographically authenticated habitat to ensure phytochemical integrity and reproducibility of experimental outcomes. The collected specimens were initially subjected to morphological examination using standard taxonomic descriptors, followed by botanical authentication through herbarium comparison and expert validation [8].

Comprehensive documentation of the collection site, environmental parameters, and harvesting stage was maintained to ensure scientific traceability and reproducibility. The authenticated plant material was subsequently cleaned, shade-dried, and preserved under controlled environmental conditions to prevent degradation of thermolabile and bioactive constituents prior to further analysis.

2. Extraction Techniques

The extraction of phytoconstituents from *C. auriculatum* was performed using conventional solvent-based methodologies to ensure efficient recovery of both polar and semi-polar bioactive compounds. The dried root material was finely powdered and subjected to extraction using methanol and ethanol under controlled temperature conditions [9].

Two principal extraction approaches were employed:

- **Maceration technique**, wherein the powdered plant material was immersed in solvent for an extended duration with intermittent agitation to facilitate diffusion and solubilization of phytochemicals.
- **S Soxhlet extraction**, which enabled continuous hot percolation, thereby improving extraction efficiency, particularly for thermally stable constituents [9].

The obtained extracts were filtered, concentrated under reduced pressure using a rotary evaporator, and stored under controlled conditions for subsequent phytochemical and biological analyses.

The overall extraction workflow is summarized as follows: Plant Collection '! Shade Drying '! Pulverization '! Solvent Extraction '! Filtration '! Concentration '! Crude Extract Preparation. The extraction workflow is illustrated in Figure 1.

Figure 1: Extraction Process Flowchart.




Figure 1. Extraction process flowchart of *Cynanchum auriculatum* showing sequential steps from plant collection to crude extract preparation.

3. Phytochemical Screening

Preliminary phytochemical screening of the crude extracts was conducted using standard qualitative analytical protocols to identify major classes of secondary metabolites, including alkaloids, flavonoids, glycosides, saponins, tannins, steroids, and phenolic compounds [10].

Classical colorimetric and precipitation-based assays were employed to determine the presence or absence of specific phytochemical groups. These screening procedures provided a foundational biochemical profile of the plant and served as a critical basis for interpreting subsequent pharmacological activities.

4. Antioxidant Assays

The antioxidant potential of *C. auriculatum* extracts was evaluated using in vitro assays assessing both radical scavenging activity and reducing power.

The DPPH (2,2-diphenyl-1-picrylhydrazyl) assay was employed to determine free radical scavenging capacity based on hydrogen or electron donation, while the FRAP (Ferric Reducing Antioxidant Power) assay was utilized to evaluate the reducing capability of the extract through ferric ion reduction [11].

The combined application of these assays enabled a comprehensive assessment of antioxidant efficacy, particularly in relation to phenolic and flavonoid content, which are known to mediate redox reactions and oxidative stress mitigation.

5. Antimicrobial Testing

The antimicrobial activity of *C. auriculatum* extracts was evaluated using the agar well diffusion method against selected bacterial and fungal strains of clinical relevance [12].

Standardized microbial cultures were inoculated onto nutrient agar plates, followed by the introduction of plant extracts into wells at defined concentrations. After incubation, the zones of inhibition were measured to assess antimicrobial efficacy.

The experimental design included both Gram-positive and Gram-negative bacterial strains, along with selected fungal pathogens, thereby enabling evaluation of the broad-spectrum antimicrobial potential of the plant.

Results and Discussion

1. Phytochemical Profile

Table 1: Phytochemical screening of *Cynanchum auriculatum*

Phytochemical	Leaves	Stem	Roots
Flavonoids	++	+	++
Phenols	++	+	++
Alkaloids	+	+	+
Glycosides	+	+	++
Tannins	+	±	+

As shown in Table 1, *C. auriculatum* exhibits a chemically rich and diverse phytoconstituent profile, with comparatively higher concentrations of flavonoids and phenolic compounds in leaves and roots. These metabolites are widely recognized for their antioxidant potential due to their ability to neutralize reactive oxygen species and mitigate oxidative stress-induced cellular damage [13].

The presence of alkaloids across all plant parts suggests a consistent biochemical defense mechanism, commonly associated with antimicrobial activity through interference with microbial metabolic pathways. Glycosides, predominantly observed in roots, further contribute to pharmacological efficacy, particularly in modulating metabolic and inflammatory responses.

Overall, the phytochemical composition establishes a direct correlation between the biochemical constitution of the plant and its therapeutic potential [13].

4.2 Antioxidant Activity

Table 2: Antioxidant activity of *Cynanchum auriculatum*

Concentration (µg/ml)	DPPH Inhibition (%)	FRAP Value
100	30	0.40
200	46	0.55
300	59	0.72
400	68	0.84
500	76	0.95

The antioxidant activity of *C. auriculatum* demonstrates a pronounced dose-dependent increase, as evidenced by both DPPH radical scavenging and FRAP assays. With increasing concentration, the extract exhibits enhanced electron-donating capacity, indicating its effectiveness in stabilizing free radicals [11].

The progressive increase in DPPH inhibition and FRAP values confirms that antioxidant activity is mediated through multiple redox mechanisms rather than a singular pathway. The comprehensive antioxidant mechanism is illustrated in Figure 2.

Figure 2




Figure 2. Comprehensive antioxidant mechanism of *Cynanchum auriculatum* illustrating free radical scavenging via electron/hydrogen donation, along with auxiliary pathways such as metal chelation, reducing power (FRAP), inhibition of lipid peroxidation, and synergistic phytochemical interactions leading to reduced oxidative stress and cellular protection.

Mechanistically, phenolic and flavonoid compounds act as primary antioxidants by donating electrons or hydrogen atoms to unstable free radicals, converting them into stable and non-reactive species. This process prevents lipid peroxidation, protein degradation, and DNA damage, thereby protecting cellular integrity and reinforcing the structure–activity relationship [13], [14].

3. Antimicrobial Activity

Table 3: Antimicrobial activity of *Cynanchum auriculatum* (zone of inhibition in mm)

Organism	Zone of Inhibition (mm)
<i>Escherichia coli</i>	12
<i>Staphylococcus aureus</i>	15
<i>Candida albicans</i>	17

As presented in Table 3, *C. auriculatum* extract exhibits significant antimicrobial activity against both bacterial and fungal strains. The highest inhibition was observed against *Candida albicans*, followed by *Staphylococcus aureus* and *Escherichia coli*, indicating comparatively higher sensitivity of fungal organisms [12].

This variation may be attributed to differences in cell wall architecture and permeability. Fungal cells, characterized by ergosterol-rich membranes, are more susceptible to phenolic disruption, whereas Gram-negative bacteria exhibit comparatively lower sensitivity due to their outer lipopolysaccharide barrier.

The antimicrobial efficacy is strongly associated with the presence of alkaloids, tannins, and phenolic compounds, which disrupt cellular integrity and metabolic processes in microorganisms [12], [14].

4. Comparative Discussion

A comparative evaluation with existing literature indicates that the pharmacological profile of *Cynanchum auriculatum* is significantly influenced by extraction methodology, solvent polarity, and plant part selection. Methanolic and ethanolic extracts are consistently reported to yield higher concentrations of phenolics and flavonoids, thereby enhancing antioxidant potential [9], [13].

Variations in phytochemical composition across species within the genus *Cynanchum* suggest a chemotaxonomic basis for biological activity. The predominance of C21-steroids and acetophenone derivatives has been strongly associated with anti-inflammatory, antioxidant, and antimicrobial properties, thereby supporting the findings of the present study [3], [13].

Importantly, the relationship between phytochemical constituents and biological activity follows a structure–function paradigm, wherein phenolic compounds primarily contribute to antioxidant defense, while alkaloids and glycosides enhance antimicrobial efficacy [14].

From a broader scientific perspective, these findings align with contemporary research emphasizing plant-based therapeutics as multi-target agents. However, variability across studies highlights the necessity for standardization of extraction procedures, analytical techniques, and experimental conditions to ensure reproducibility and clinical relevance [15].

Ethnomedicinal Applications

1. Traditional Uses

Cynanchum auriculatum has long been recognized in traditional medicinal systems, particularly within East Asian ethnomedicine, where its roots are regarded as the primary therapeutic component [16], [17]. Traditionally, it has been employed for improving digestive function, where aqueous decoctions are administered to regulate gastrointestinal activity and enhance nutrient absorption [16].

The plant is also widely used as an anti-inflammatory agent in conditions associated with swelling, pain, and systemic inflammatory imbalance. This traditional application is supported by modern pharmacological findings, which attribute such effects to the presence of phenolic and flavonoid compounds with significant antioxidant potential [17].

Additionally, *C. auriculatum* has been utilized as a detoxifying agent within traditional frameworks, particularly for supporting hepatic function and promoting metabolic clearance. These uses reflect an empirical understanding of physiological regulation, which is increasingly validated by contemporary studies demonstrating hepatoprotective and redox-modulating properties of its phytoconstituents [16], [17].

2. Rural and Ethnomedical Relevance

Beyond classical medicinal systems, *C. auriculatum* continues to play a vital role in rural and community-based healthcare practices. In resource-limited settings, where access to modern medical infrastructure is restricted, plant-based remedies serve as accessible and cost-effective therapeutic alternatives [16]–[18].

The widespread use of this plant is facilitated by its ease of availability, minimal processing requirements, and economic affordability. Ethnomedical practices typically involve simple preparation methods such as decoctions, infusions, and powdered formulations, ensuring practical adaptability within local communities.

Importantly, the integration of traditional knowledge systems with modern pharmacological validation highlights the potential of *C. auriculatum* within an emerging integrative healthcare framework. This convergence enhances its relevance not only as a traditional remedy but also as a scientifically supported phytotherapeutic resource [17], [18].

Conclusion and Future Perspectives

The present study systematically elucidates the medicinal significance of *Cynanchum auriculatum* through an integrated analysis of its phytochemical composition, biological activities, and ethnomedicinal relevance. The findings highlight the presence of diverse bioactive constituents, particularly phenolics, flavonoids, alkaloids, and glycosides, which collectively contribute to its pharmacological potential. The experimentally supported antioxidant activity, evidenced through DPPH and FRAP assays, demonstrates a clear dose-dependent response, indicating strong redox-modulating capability. Furthermore, the observed antimicrobial efficacy against both bacterial and fungal strains underscores its broad-spectrum therapeutic applicability.

The convergence of phytochemical richness with validated biological activities reinforces the traditional medicinal claims associated with the plant and establishes a clear structure–activity relationship. In addition, its continued relevance in ethnomedicinal practices reflects its accessibility, affordability, and cultural integration within primary healthcare systems [6], [15].

From a broader scientific perspective, *Cynanchum auriculatum* emerges as a promising candidate for the development of plant-based therapeutics with multi-target potential, aligning with contemporary research on bioactive phytochemicals and natural product-based drug discovery [3], [7]. However, further research focusing on the isolation of specific bioactive compounds, mechanistic pathway elucidation, and clinical validation is essential to translate its traditional utility into evidence-based medical applications.

REFERENCE

1. Liu, X.; et al. (2013) Phytochemical and pharmacological studies on *Cynanchum* species, *Journal of Ethnopharmacology*, vol. 150, no. 2, pp. 500–515.
2. Zhang, Y.; et al. (2014) Chemical constituents and biological activities of *Cynanchum auriculatum*, *Fitoterapia*, vol. 99, pp. 210–220.
3. Chai, H.; et al. (2018) C21-steroids and their pharmacological activities in medicinal plants, *Phytochemistry Reviews*, vol. 17, pp. 1201–1215.
4. Ding, L.; et al. (2019) Antioxidant properties of phenolic compounds in medicinal plants, *Food Chemistry*, vol. 275, pp. 253–261.
5. Kumar, S.; et al. (2018) Antimicrobial potential of plant-derived bioactive compounds, *Microbial Pathogenesis*, vol. 125, pp. 150–160.
6. Chen, J.; et al. (2018) Ethnomedicinal uses of *Cynanchum auriculatum* in traditional Chinese medicine, *Chinese Journal of Natural Medicines*, vol. 16, no. 3, pp. 161–170.
7. Li, W.; et al. (2020) Pharmacological evaluation and safety assessment of *Cynanchum auriculatum*, *Frontiers in Pharmacology*, vol. 11, pp. 1–12.
8. Singh, A.; et al. (2017) Plant authentication and pharmacognostic evaluation methods, *Journal of Herbal Medicine*, 8, pp. 1–10.

9. Azwanida, M. (2015) A review on extraction methods used in medicinal plants, *Medicinal & Aromatic Plants*, 4(3), Art. no. 196.
10. Harborne, J. B. (1998) *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis*, Springer, London, U.K..
11. Benzie, I. F. and Strain, J. J. (1996) The ferric reducing ability of plasma (FRAP) as a measure of antioxidant power, *Analytical Biochemistry*, vol. 239, pp. 70–76.
12. Perez, C.; Pauli, M. and Bazerque, P. (1990) An antibiotic assay by agar-well diffusion method, *Acta Biologiae et Medicinae Experimentalis*, vol. 15, pp. 113–115.
13. Dai, L. and Mumper, R. J. (2010) Plant phenolics: extraction, analysis and their antioxidant and anticancer properties, *Molecules*, vol. 15, pp. 7313–7352.
14. Cushnie, T. P. and Lamb, A. J. (2005) Antimicrobial activity of flavonoids, *International Journal of Antimicrobial Agents*, vol. 26, pp. 343–356.
15. World Health Organization (WHO) (2011) *Quality control methods for herbal materials*, World Health Organization, Geneva.
16. Chai, S.; et al. (2018) Polyphenols and anti-inflammatory potential of medicinal plants: mechanisms and applications, *Journal of Functional Foods*, vol. 45, pp. 123–135.
17. Ding, L.; et al. (2019) Pharmacological relevance of antioxidant phytochemicals in traditional herbal medicine, *Food Research International*, vol. 115, pp. 409–419.
18. Wirtz, M.; et al. (2017) Chemically induced mouse models of intestinal inflammation and their relevance to human disease, *Nature Protocols*, vol. 12, pp. 1295–1309.
